

# WESTCHASE GASTROENTEROLOGY

JOHN CHANG, MD, FACG   AMIR AWAD, MD, FACG   ALFREDO MENDOZA, MD, MS

4695 Van Dyke Road, Lutz FL 33558 || 11912 Sheldon Road, Tampa FL 33626 || 508 S. Habana Ave, Ste 270, Tampa FL 33609

Telephone: 813.920.8882   Fax: 813.920.8883

[www.westchasegi.com](http://www.westchasegi.com)

## PREPOPIK Day-Before Dosing Instructions

Please read over these instructions carefully

**Day of your procedure:** \_\_\_\_\_ . Be at the Outpatient Endoscopy Unit at: \_\_\_\_\_ .

- ✓ **DO NOT eat or drink.** Exception: if you take heart, seizure, blood pressure and/or thyroid medications in the morning, take with a small sip of water at least two (2) hours before your procedure time.
- ✓ Please bring the following items with you:
  1. Your insurance cards and driver's license
  2. Eye glasses if needed for reading
- ✓ Please leave jewelry and valuables at home, wedding band(s) may be worn.
- ✓ Please wear comfortable clothes; you will be changing into a gown.

### Important reminders:

- ✓ Someone must drive you home after the procedure. You will not be discharged from the facility if you do not have a driver.
- ✓ Plan on resting after your procedure. Avoid big activities, operating motor vehicles, and do not drink alcohol on the day of the procedure.
- ✓ If you are DIABETIC:
  - a. If you use insulin only **use ½ dose** of insulin the night before your procedure and **NO** insulin the morning of your procedure. You may resume your normal dose after your procedure.
  - b. If you take pills, **DO NOT** take the day before and the morning of your procedure. You may resume your normal dose after your procedure.
- ✓ **DO NOT take for SEVEN (7) days prior** to procedure: any **anti-coagulants/blood thinners/anti-platelets** medication(s) (Plavix, Coumadin, Effient, Xarelto, Pradaxa, etc.) Aspirins, and/or NSAIDs. **Please contact in advance the physician who prescribed any of those medications for clearance and/or instructions.** This includes over the counter (OTCs): iron, multi-vitamin with iron, Vitamin E, St John's Wort, ginkgo-biloba, ginger, fish oil, and garlic pills. *If you need to take a pain reliever, you can take Tylenol (Acetaminophen).*
- ✓ Reduce fiber intake for 2 to 3 days prior to procedure. For example, do not eat nuts, seeds, popcorn, corn, etc. Do not use fiber supplements such as Metamucil, Citrucel, Fiberall, etc.
- ✓ **Please fill your laxative prep prescription at least three (3) days in advance to your procedure to ensure the availability of the medication at your pharmacy or to avoid any unforeseen circumstances.**

NOTE: IF YOU HAVE NOT SCHEDULED A FOLLOW-UP APPOINTMENT FOR AFTER YOUR PROCEDURE, PLEASE CALL THE OFFICE TO SCHEDULE ONE. WE **DO NOT** GIVE RESULTS OVER THE PHONE, NO EXCEPTIONS. THANK YOU! **IF YOU HAVE ANY QUESTIONS, PLEASE CALL THE OFFICE.**

## **THE DAY BEFORE YOUR PROCEDURE: CLEAR LIQUID DIET ALL DAY**

Please **DO NOT** eat breakfast or **ANY** solid food the day before and the morning of your procedure. For example, if your procedure is on Monday, do not eat breakfast or any solid food all day Sunday and Monday morning. You may resume regular diet as tolerated after your procedure.

**Drink at least 8 ounces of clear liquids every hour up until you start the PREPOPIK**

**PREPOPIK preparation consists of two (2) regimens taken with a clear liquid of your choice.**

|  |   |
|--|---|
| <p><u>First regimen: take @ 4 PM</u></p> <p>A. Follow package directions for how to mix the powder pack. Take as instructed.</p> <p>Drink five (5) eight-ounce (8 oz) of clear liquids before your second dose of PREPOPIK</p> | <p><u>Second regimen: take @ 10 PM</u></p> <p>B. Repeat step A</p> <p>Drink three (3) eight-ounce(8 oz) of clear liquids before bedtime</p> |
|--|---|

There has to be a six (6) hour time span between each regimen

**One hour** after completing PREPOPIK second regimen: take by mouth four (4) tablets of Dulcolax (aka Bisacodyl) 5mg/tab.

Please stay well hydrated before, during, and after you have completed your Prepopik, **HOWEVER**, you must **stop all liquids intake four (4) hours prior to your procedure time, this also means coffee.** (For example: if your procedure time is 8 am, do not drink after 4 am.)

**IMPORTANT REMINDER: DO NOT EAT OR DRINK ANY: DAIRY & NON DAIRY FOODS; RED, PURPLE, & BLUE COLORED LIQUIDS; JUICES WITH PULP/UNSTRAINED FRUIT**

| Food Groups                     | Recommended  | Avoid   |
|---------------------------------|--|---|
| <b>Milk &amp; milk products</b> | none   | all   |
| <b>Vegetables</b>               | none   | all   |
| <b>Fruits</b>                   | fruit juices without pulp  | nectars; all fresh, canned, and frozen fruits |
| <b>Breads &amp; grains</b>      | none   | all   |
| <b>Meat or meat substitutes</b> | none   | all   |
| <b>Fats &amp; oils</b>          | none   | all   |
| <b>Sweets &amp; desserts</b>    | gelatin, fruit ice, popsicle without pulp, clear hard candy  | all others                                    |
| <b>Beverages</b>                | black coffee only; tea; soft drinks; water; lactose-free, low residue supplements if approved by physician | all others                                    |
| <b>Soups</b>                    | bouillon, consommé fat free broth  | all others                                    |